



# Viscosity solutions for realistic Shape-From-Shading

Emmanuel Prados, Olivier Faugeras

## ► To cite this version:

Emmanuel Prados, Olivier Faugeras. Viscosity solutions for realistic Shape-From-Shading. Workshop on Numerical Methods for Viscosity Solutions and Applications, Sep 2004, Rome, Italy. inria-00590177

**HAL Id: inria-00590177**

**<https://inria.hal.science/inria-00590177>**

Submitted on 6 May 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


# Viscosity solutions For Realistic Shape-From-Shading

**Emmanuel Prados & Olivier Faugeras**  
Odysée Lab., INRIA Sophia Antipolis, France



**Workshop “Numerical Methods for Viscosity Solutions and Applications”**  
**Rome, September 6-8, 2004**

# The Shape From Shading Problem



## Problem:

- Inverse problem to image synthesis,
- Recover the surface(s) which yield(s) the same image.

## Used Information:

- Shading.

# Shape From Shading Data

- One image;
- Intrinsic parameters of the camera:
  - Focal length,
  - Size of the pixels
- Surface parameters:
  - Reflectance of the surface (albedo)
- Lighting parameters:
  - Flash...
- *Database of real images of faces available online:*
  - Experimental setup: scene illuminated by the flash of the camera.
  - Parameters are detailed.
  - WWW link:



<http://www-sop.inria.fr/odyssee/team/Emmanuel.Prados/>

# Report:

- 1) Most of the SFS methods model the problem very basically:  
Ex.: Camera=Orthographic Projection...  
⇒ Not Realistic!
- 2) Lots of SFS algorithms require boundary data (generally Dirichlet BC)  
⇒ Not Available with **Real Images!**



The results on real images are generally very disappointing.

# Shape From Shading Contributions:

- We propose **more realistic modeling**:  
Example: camera = pinhole  
(perspective projection)
- We **reduce as much as possible** the  
requirement of **the boundary data**  
(in particular on the boundary of the images).

# Modeling / Mathematical formulations

## ■ The Basic Assumption: Lambertian Surface


(without specularities or highlights)

⇒ Involves a connection between:

- the brightness of the image (intensity) : variable ***I***,
- the normal surface: ***n***, <<Depends on the parameterization>>.
- the lighting direction: ***L*** <<Depends on the lighting>>.

$$I(X) = \cos(\mathbf{n}(X), \mathbf{L}(X)) = \frac{\mathbf{n}(X)}{|\mathbf{n}(X)|} \cdot \mathbf{L}(X)$$

# Modeling / Mathematical formulations



## ■ Orthographic projection (classical modeling)

- Far light source:

$$I(x)\sqrt{1 + |\nabla u|^2} + \nabla u \cdot \mathbf{l} - \gamma = 0,$$

- Frontal far light source: (Eikonal equation)

$$|\nabla u| = f(x),$$

$$f(x) = \sqrt{\frac{1}{I(x)^2} - 1}.$$

## ■ More realistic Modeling: Perspective Projection:


- Far light source:

$$I(x)\sqrt{f^2|\nabla u|^2 + (x \cdot \nabla u + 1)^2} - (f\mathbf{l} + \gamma x) \cdot \nabla u - \gamma = 0,$$

- Light source located at the Optical center:

$$I(x)\sqrt{f^2|\nabla u|^2 + (x \cdot \nabla u)^2 + Q(x)^2} - Q(x) = 0.$$

$$Q(x) = \frac{f}{\sqrt{|x|^2 + f^2}}.$$





# A Generic Hamiltonian

- All these SFS equations are special cases of the “generic” equation:

$$\kappa_x \sqrt{|A_x \nabla \mathbf{u} + \mathbf{v}_x|^2 + K_x^2} + \mathbf{w}_x \cdot \nabla \mathbf{u} + c_x = 0 \quad (1)$$

Where  $\kappa_x, K_x \geq 0$ ,  $A_x \in \mathcal{M}_{2 \times 2}$ ,  
 $\mu_x, \nu_x \in \mathbb{R}^*$ ,  $\mathbf{v}_x, \mathbf{w}_x \in \mathbb{R}^2$  and  $c_x \in \mathbb{R}$ .

(1) is a HJ PDE of the form  $H(x, \nabla u)$  with:

$$H(x, \mathbf{p}) = \kappa_x \sqrt{|A_x \mathbf{p} + \mathbf{v}_x|^2 + K_x^2} + \mathbf{w}_x \cdot \mathbf{p} + c_x.$$

$H(x, \mathbf{p})$  convex in  $\mathbf{p}$ .

# Ill-posedness of the SFS problem

- SFS equations are of the form:

$$H(x, \nabla u) = 0, \quad x \text{ in } \Omega \quad (\text{a bounded set}).$$


⇒ They require **Boundary Conditions** (BC) on  $\partial\Omega$

⇒ But the BC on  $\partial\Omega$  are **not sufficient!**

**SFS equations can have singularities:**

*For all the previous SFS equations,  
the set of singular points is*

$$\{ x \mid I(x) \text{ is maximal (i.e 1)} \}$$





# Recall:

## In Shape from Shading

### **Data = one image.**

- ⇒ We do not have boundary data;
- ⇒ i.e. We do not know what are the adequate boundary conditions.

# Solution proposed in the SFS literature

Notion	SFS Biblio.	Data (DBC)	Limitations
<b>Continuous</b> Viscosity solutions (Crandall-Lions...)	Lions/Rouy/ Tourin (92)	$S_{ing} \cup \partial\Omega$	<ul style="list-style-type: none"> <li>Compatibility conditions.</li> <li>Dirich. data on <math>S_{ing} \cup \partial\Omega</math></li> </ul>
$C^1$ Functions	Dupuis/ Oliensis (94)	Set of local $Min.$ of $u$	Value function ? <b>Few intuitive...</b> But: <b>Interest :</b> <b>Minimal Data are required!</b>
<b>Discontinuous</b> Viscosity solutions (Ishii...)	Prados/ Faugeras (03)	$N(S_{ing}) \cup \partial\Omega$	<ul style="list-style-type: none"> <li>Dirichlet data on <math>N(S)_{ing} \cup \partial\Omega</math></li> </ul>
<b>Singular</b> Viscosity solutions (Camilli-Siconolfi:99)	Camilli/ Falcone (96)	$\partial\Omega$	<ul style="list-style-type: none"> <li>not always adapted to SFS.</li> <li>still requires data on <math>\partial\Omega</math></li> </ul>

# Contributions:

- We have slightly extended and modified the notion of singular viscosity solutions (Camilli/Siconolfi:99,02) in order to:
  - to extend the Dupuis/Oliensis' characterization of the  $C^1$  solutions to the viscosity solutions  $\Rightarrow$  data are minimal.
  - such that compatibility conditions are not required.

# Notion of SDVS

- Definition: We define a notion of solution
  - which extends the classical notions of
    - of the discontinuous viscosity solutions (Ishii...)
    - of the singular viscosity solutions (Camilli-Siconolfi:99,02).
  - Which allows to fix the values of the solutions at the singular points or on the boundary  $\partial\Omega$  when we know them, and to send them at the infinity (state constraints), when we do not know them.

# Properties of the SDVS:

- Existence of the SDVS
    - Dynamical programming principle,
    - value function;
  - Strong uniqueness;
  - Stability results;
- ➔ Application to the Shape From Shading problem...

# Original theorems:

- Theorem: (roughly)  
« A discontinuous viscosity solution without “local minima” is the SDVS».
- Corollary:  
Characterization of the discontinuous viscosity solutions by their « minima ».



# Numerical approximations:

- 4 stages:

- i) Management of the state constraints;

- ii) Regularization of the equations

- In SFS (we truncate the image,  $I_\varepsilon = \min(I(x), 1 - \varepsilon)$ );


- iii) Designing of the approximation scheme;

- iv) Deduction and implementation of the associated algorithm.

# A Numerical Result on a Synthetic Image (Mozart)




a) Synthetic image of the  
classical Mozart face




b) Reconstruction

# Numerical results on Real images (Faces)...




a) Real image of a face




b) Reconstruction

# Numerical results on Real images (Stomach)...




a) Real image of a stomach




b) Reconstruction

# Numerical results on Real images (Pages)



a) Real image of a page



b) Reconstruction

# Conclusion:

- SFS problem  $\neq$  Eikonal equation.
  - Realistic SFS consists in solving more complex equations: for ex. the “generic SFS” equation.
- In SFS, we do not have at our disposal boundary data.
  - The notion of SDVS proposes some more adapted solutions than the other classical solutions.

# For more details:

## ■ Theoretical part:

See the INRIA research report:

"A viscosity method for Shape From Shading without boundary data"  
(*E. Prados, F. Camilli and O. Faugeras*), August 2004.

## ■ Numerical part: See

- the INRIA research report No RR-5005 :

"A mathematical and algorithmic study of the Lambertian SFS  
problem for orthographic and pinhole cameras"  
(*E. Prados and O. Faugeras*), September 2003.

- E. Prados' PhD thesis, October 2004.

## ■ Web page:

<http://www-sop.inria.fr/odyssee/team/Emmanuel.Prados/>

## ■ Thanks!